

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Appl. No. : 09/806,370 Confirmation No. 8568
Applicant : Randall K. Holmes et al
Filed : October 3, 2001
TC/A.U. : 1645
Examiner : Virginia A. Portner
Docket No. : 33,383-00
Customer No. : 38199

Mail Stop RCE
Commissioner for Patents
P. O. Box 1450
Alexandria, VA 22313-1450

DECLARATION

Sir:

I, Mary E. Bak, residing at 1415 Comly Court, Maple Glen, PA, 19002, a citizen of the United States of America, do declare and state that:

1. I am one of the named attorneys of record in the above-identified patent application.
2. This Declaration is submitted in the above-identified application in response to the Examiner's rejection under 35 USC § 112, first paragraph in the Office Action dated March 10, 2004.

Express Mail No. EU531571678US

BEST AVAILABLE COPY

3. Specifically, this Declaration is submitted to support the insertion by amendment of SEQ ID NO: 1 into the specification, which is the mature wild-type cholera holotoxin subunit A sequence as set forth in Domenighini *et al.*, International Patent Publication No. WO 93/13202 (hereinafter Domenighini) cited in the specification at page 38, line 10 and properly incorporated by reference. See, Exhibit A attached herewith.

4. The same sequence appears in Mekalanos *et al.*, 1983 *Nature*, 306:551-557 (hereinafter Mekalanos) cited in the specification at page 2, line 4 in the context of the entire CT sequence with subunit B and 5' and 3' untranslated regions. See, Exhibit B. The mature subunit A is indicated in Mekalanos by the first amino acid appearing under the first mature amino acid "N" in the sequence. SEQ ID NO: 1 is a duplicate of the mature subunit A sequence as set forth in both Domenighini and Mekalanos.

5. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Date: July 12, 2004

By: Mary E. Bak
Mary E. Bak



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C12N 15/00, 15/31, A61K 39/106 A61K 39/108, C12P 21/02 C12N 1/21 // (C12N 1/21 C12R 1:19)	A1	(11) International Publication Number: WO 93/13202 (43) International Publication Date: 8 July 1993 (08.07.93)
--	----	---

(21) International Application Number: PCT/EP92/03016

(22) International Filing Date: 30 December 1992 (30.12.92)

(30) Priority data:
MI91A03513 31 December 1991 (31.12.91) IT(71) Applicant (for all designated States except US): BIOGINE
SCLAVO SPA [IT/IT]; Via Fiorentina, 1, I-53100 Siena
(IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DOMENIGHINI, Mar-
io [IT/IT]; Via Ungaretti, 17, I-53010 Quercegrossa (IT).
RAPPUOLI, Rino [IT/IT]; Via Calamandrei, 39, I-
53010 Quercegrossa (IT). PIZZA, Mariagrazia [IT/IT];
Via Colombini, 30, I-43100 Siena (IT). HOL, Wim [NL/
US]; 18332 57th Avenue, N.E., Seattle, WA 98155 (US).(74) Agent: HALLYBONE, Huw, George; Carpmals & Rans-
ford, 43 Bloomsbury Square, London WC1A 2RA (GB).(81) Designated States: AT, AU, BB, BG, BR, CA, CH, CS,
DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG,
MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, US,
European patent (AT, BE, CH, DE, DK, ES, FR, GB,
GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF,
BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG).


Published

With international search report.

Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.(54) Title: IMMUNOGENIC DETOXIFIED MUTANTS OF CHOLERA TOXIN AND OF THE TOXIN LT, THEIR PRE-
PARATION AND THEIR USE FOR THE PREPARATION OF VACCINES

(57) Abstract

An immunogenic detoxified protein comprising the amino acid sequence of subunit A of cholera toxin (CT-A) or subunit A of an *Escherichia coli* heat labile toxin (LT-A) or a fragment thereof wherein one or more amino acids at, or in positions corresponding to Val-53, Ser-63, Val-97, Tyr-104 or Pro-106 are replaced with another amino acid or deleted. Examples of specific replacements include Val-53-Asp, Val-53-Glu, Val-53-Tyr, Ser-63-Lys, Val-97-Lys, Val-97-Tyr, Tyr-104-Lys, Tyr-104-Asp, Tyr-104-Ser, Pro-106-Ser. The immunogenic detoxified protein is useful as vaccine for *Vibrio cholerae* or an enterotoxigenic strain of *Escherichia coli* and is produced by recombinant DNA means by site-directed mutagenesis.



LT2	1	--.FF-----T---R-A---L---QQ-AYE---PI---	38
LT1	1	-----FRS-----	39
LT1_1A	1	-G-R-----R-----HN-----	40
CT	1	NDDKLYRADSRPPDEIKQSGGLMPRGQSEYFDRGTQMNN	40
		-----E-----V--NT--N-----TVT--Q---I--N--GS-	78
		-----Y-----	79
		-----Y-----L-----A--S-----Y	80
		LYDHARGTQTGFVRHDDGYVST ¹ SISLRSAHLVGQTILSGH	80
		NE-----V-P---L-D--G---R---Y-S-N-FA-----	118
		-LTIYI---...-----IS-----	116
		-----V-----Y-----	120
		STVYIYVIATAPNMFNVNDVLGAY ¹ SPHPDEQEVSALGGIP	120
		L---I-----SF-A-EGGMQ---D--GDLF-G-TV--N--	158
		-----	156
		-----N---I--R-----E-----R--N---E-	160
		YSQIYGWYRVHFGVLDEQLHRNRGYRDRYYSNLDIAPAAD	160
		--Q-----SNFP---M--STF--EQ-VPNNKEFK-GV-I	198
		-----	196
		--R-----D-Q-----Q---DSS-TITGD--N	200
		GYGLAGFPPEHRAWREEPWIIHAPPGCGNAP ¹ SSMSNTCD	200
		SA-NV--KYD-MNFKKLL--RLALTFFM--D-F-GVHGE----	241
		-----	236
		-E--N-STIY-R-----D-.--EV-.IY---.R---	240
		EKTQSLGVKFLDEYQSKVKRQIFSGY.QSDID.THNRI.KDEL	240

Figure 1

Cholera toxin genes: nucleotide sequence, deletion analysis and vaccine development

John J. Mekalanos, Daryl J. Swartz & Gregory D. N. Pearson

Department of Microbiology and Molecular Genetics, Harvard Medical School, 25 Shattuck Street, Boston, Massachusetts 02115, USA

Nigel Harford, Françoise Groyne & Michel de Wilde

Department of Molecular Genetics, Smith-Kline-R.I.T., rue de l'Institut, 89, B-1330 Rixensart, Belgium

Nucleotide sequence and deletion analysis have been used to identify the regulatory and coding sequences comprising the cholera toxin operon (ctx). Incorporation of defined in vitro-generated ctx deletion mutations into Vibrio cholerae by in vivo genetic recombination produced strains which have practical value in cholera vaccine development.

MODERN history has recorded seven world pandemics of cholera, a diarrhoeal disease produced by the Gram-negative bacterium *Vibrio cholerae*¹. Laboratory tests can distinguish two biotypes of *V. cholerae*, classical and El Tor, the latter being responsible for the most recent cholera pandemic. The diarrhoeal syndrome induced by colonization of the human small bowel by either biotype of *V. cholerae* is caused by the action of cholera toxin, a heat-labile enterotoxin secreted by the growing vibrios². Diarrhoeal diseases affecting both humans and animals caused by some enterotoxinogenic strains of *Escherichia coli* are also induced by a heat-labile enterotoxin (LT) which is closely related to cholera toxin in structure and mode of action¹⁻³.

Cholera toxin is an 84,000-molecular weight (MW) protein composed of one A subunit (27,000 MW) and five B subunits (11,600 MW). The A subunit, although synthesized as a single polypeptide chain, is usually proteolytically nicked to form two disulphide-linked polypeptides, A1 (22,000 MW) and A2 (5,000 MW)^{4,5}. The A1 polypeptide is an enzyme and promotes the activation of adenylate cyclase in target cells by catalysing the ADP-ribosylation of a GTP-binding regulatory component of the cyclase complex⁶. The resulting accumulation of cyclic AMP in the intestinal mucosa leads to the severe fluid loss characteristic of cholera. Each B subunit has a high binding affinity for the toxin's cell surface receptor, ganglioside GM₁ (ref. 7). Neutralizing antibodies raised against the holotoxin react mainly with the B subunits¹⁻².

Much of the current interest in the genetics of cholera toxin has been promoted by the need to develop a more efficacious vaccine against this enterotoxic disease. Parenterally administered, killed whole-cell and toxoid vaccines have been shown to be largely ineffective in producing long-lasting immunity to cholera, presumably because they lack the ability to induce local immune responses in the intestine^{8,9}. Since the natural disease is capable of inducing prolonged immunity^{9,10}, several investigators have proposed the use of attenuated, non-toxinogenic mutants of *V. cholerae* as live oral cholera vaccines¹¹⁻¹⁶. While encouraging results in volunteer studies have been obtained with some of these strains, factors such as genetic instability or poor colonizing ability have contraindicated their use in the field¹⁵⁻¹⁷.

The recent relaxation of US governmental guidelines prohibiting the molecular cloning of bacterial toxin genes has permitted the use of a powerful new approach to the analysis of cholera toxin gene structure and vaccine development. These studies have shown that like the *elt* genes, which encode *E. coli* LT¹⁸, the genes for the A and B subunits of cholera toxin are arranged in a single transcriptional unit with the A cistron (*ctxA*) preceding the B cistron (*ctxB*)¹⁹. *V. cholerae* strains of the classical biotype contain a nontandem, chromosomal duplication of the *ctx* operon that is structurally identical in all strains.²⁰ In contrast, about 70% of El Tor strains have only a single copy of

ctx, while the remaining strains have two or more *ctx* copies present on a tandemly repeated genetic element. This genetic duplication and amplification of the toxin operon may be related to the instability observed in some of the earlier *V. cholerae* toxin mutants^{13,16}.

In this article, we report the entire nucleotide sequence of one *ctx* operon together with partial sequences containing the *ctx* promoter regions of five other cloned *ctx* copies. Deletion analysis has allowed the identification of toxin transcriptional and translational regulatory sequences. An *in vitro*-constructed, internal deletion in *ctxA* was recombined *in vivo* into both *ctxA* gene copies of *V. cholerae* strain Ogawa 395. Since this genetic recombinant still produces the immunogenic B subunit of the toxin, it should have practical value in cholera vaccine development.

Molecular cloning of *ctx* operon copies

A total of six *ctx* copies were cloned from four *V. cholerae* strains. These include both *ctx* copies from strain 569B, both copies from strain RV79, one of two copies from strain E7946 and the single *ctx* copy of strain 2125. With the exception of the classical strain 569B, all these strains are El Tor in biotype. The *ctx* copies were cloned as various *V. cholerae* restriction fragments which hybridized with ³²P-labelled *elt* or *ctx* probes as described in the legend to Fig. 1.


The restriction sites for several endonucleases were located on these cloned inserts, and the resulting maps were aligned at the conserved *Xba*I site previously determined to lie early in the A cistron¹⁹, (Fig. 1A). Other conserved restriction sites for *Nru*I, *Pst*I, *Ava*I and *Bgl*II were also found preceding *ctxA* on these various inserts (Fig. 1). Additional restriction mapping and hybridization analysis has indicated that the 5 kilobase pairs (kbp) of DNA directly preceding the toxin structural genes is the same for all cloned *ctx* copies thus far examined²¹. Since the larger chromosomal sequence environment flanking these different *ctx* copies appears to vary as determined by Southern blot hybridization, we have proposed that the conserved DNA immediately upstream of *ctx* is part of a genetic element responsible for toxin operon duplication and transposition events. From a practical point of view, the conserved 5' flanking sequences associated with *ctx* provided part of the necessary homology for efficient *in vivo* recombination of *in vitro* constructed deletion mutations into multiple copies of the toxin operon (see below).

Nucleotide sequence of *ctx*

The strategy used to determine the complete nucleotide sequence of the single *ctx* copy of strain 2125 is shown below the restriction map of the insert cloned on pRIT10824 (Fig. 1B). The 2,020 nucleotides determined are shown in Fig. 2. Comparison of the 2,125 nucleotide sequence with both the *elt* nucleotide sequence^{20,21} and known amino acid sequences of

BEST AVAILABLE COPY

Fig. 2 DNA sequence of the *V. cholerae* toxin operon from strain 2125. The antisense strand is shown from 5' to 3'. From nucleotide 427 to 1,663, the sequence is compared with the published sequence of LT genes^{20,21}; *elt* nucleotides are shown above the sequence only where they differ from the *cix* sequence except between nucleotides 810 and 830, where deletion of a T (arrowed) in the *elt* sequence creates a frameshift which is corrected by insertion of a C (arrowed) 16 bp downstream (see text). Analogous events have previously been seen after pseudoreversion of frameshift mutations⁴³ but to our knowledge, this is the first naturally occurring case described. The deduced amino acid sequence of *cixA* (nucleotide 516 to 1,289) and *cixB* (nucleotide 1,289 to 1,660) is shown and compared with that of LT. Amino acids that differ in LT are shown below the *cixA* and *cixB* cistrons (nucleotides 1,289–1,292). Sequence exhibiting dyad symmetry and potentially involved in transcription termination is indicated with divergent arrows. Features of the sequence immediately upstream of the *cixA* gene are detailed in Fig. 3.



sequence (nucleotides 1,277–1,282, Fig. 2) of the *cixA* cistron. The first two nucleotides of the *cixA* translation termination signal TGA are the last two nucleotides of the *cixB* translation initiation triplet ATG. This particular overlapping arrangement is also found several times in phage λ operons³⁰ and may be involved in translational coupling³¹ of the *cixA* and *cixB* genes. However, evidence presented below suggests that this is not the case with the *cix* operon. Where documented, translational coupling is observed between cistrons whose gene products interact in a one to one stoichiometry³¹, and in contrast, the cholera toxin molecule is composed of one A subunit and five B subunits. Moreover, *E. coli* produces stoichiometrically 7 times more cholera toxin B subunit than A subunit (data not shown). Fusion of the *cixB* gene to various *E. coli* promoters allows high expression of *cixB* in the absence of *cixA* translational initiation signals. These data suggest that translation of *cixB* relies primarily on independent initiations promoted by its own ribosome binding site.

Another experiment supports this conclusion. Our DNA sequencing analysis identified two *NdeI* sites at positions 561 and 1,337 within the *cixA* and *cixB* genes, respectively. The positions of these sites relative to the reading frames of *cixA* and *cixB* allowed us to construct a *cixA* deletion which codes for an in-frame fusion of amino acid 17 of the A subunit signal sequence to amino acid 19 of the B signal and thus maintains the normal processing site of the B signal sequence (residue

21). This genetic fusion makes B subunit expression dependent on the efficiency of the A cistron translation initiation sequences, provided the hybrid signal sequence is processed at normal efficiency. *NdeI* digestion of plasmid pGP3 followed by ligation produced such a fusion between these two sites and gave plasmid pJM3.1. Plasmid pJM3.1 produced $0.056 \mu\text{g ml}^{-1}$ of B subunit in *E. coli* MS371 while pGP3 produced $0.50 \mu\text{g ml}^{-1}$. These data suggest that the *cixB* ribosome binding site is about ninefold more efficient than the *cixA* site.

Toxin promoter regions

We determined approximately 200 base pairs of sequence upstream of the *XbaI* sites for each of the other five additional cloned copies of the *cixA* gene, cloned on plasmids pGP3, pGP4, pGP5, pGP6 and JM17. Comparison of these sequences with the corresponding region of the *cixA* gene derived from strain 2,125 indicated a perfect conservation of sequence between these copies from nucleotides 413 to 590 with one notable exception. The sequence TTTTGAT comprising nucleotides 419–425, 426–432 and 433–439 of the 2,125 sequence was found tandemly repeated 3–8 times preceding different *cixA* gene copies (Fig. 3). Figure 3 shows part of a sequencing gel autoradiograph that spans DNA carrying eight of these tandem repeats in the region adjacent to the *cixA* gene of pJM17.

To determine the position of the toxin operon promoter with respect to these repeated sequences, we used nuclease *Bal31*